

# Natural Logarithms (Sect. 7.2)

- ▶ Definition as an integral.
- ▶ The derivative and properties.
- ▶ The graph of the natural logarithm.
- ▶ Integrals involving logarithms.
- ▶ Logarithmic differentiation.

# Definition as an integral

Recall:

(a) The derivative of  $y = x^n$  is

# Definition as an integral

Recall:

(a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.

# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is

# Definition as an integral

Recall:

(a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.

(b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ ,

## Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .

# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$

# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$  is neither rational nor trigonometric function.



# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$  is neither rational nor trigonometric function. This is a new function.

# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$  is neither rational nor trigonometric function. This is a new function.

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

# Definition as an integral

Recall:

- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$  is neither rational nor trigonometric function. This is a new function.

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .

# Definition as an integral

Recall:


- (a) The derivative of  $y = x^n$  is  $y' = n x^{(n-1)}$ , for  $n$  integer.
- (b) The integral of  $y = x^n$  is  $\int x^n dx = \frac{x^{(n+1)}}{(n+1)}$ , for  $n \neq -1$ .
- (c) Case  $n = -1$ :  $\int \frac{dx}{x}$  is neither rational nor trigonometric function. This is a new function.

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .




# Definition as an integral

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .



# Definition as an integral

## Definition


The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .

## Definition

The *number e* is the number satisfying  $\ln(e) = 1$ ,




# Definition as an integral

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .



## Definition

The *number e* is the number satisfying  $\ln(e) = 1$ , that is,

$$\int_1^e \frac{dt}{t} = 1.$$


# Definition as an integral

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .



## Definition

The *number e* is the number satisfying  $\ln(e) = 1$ , that is,

$$\int_1^e \frac{dt}{t} = 1.$$

( $e = 2.718281\dots$ ).




# Definition as an integral

## Definition

The *natural logarithm* is the function

$$\ln(x) = \int_1^x \frac{dt}{t}, \quad x \in (0, \infty).$$

In particular:  $\ln(1) = 0$ .




## Definition

The *number e* is the number satisfying  $\ln(e) = 1$ , that is,

$$\int_1^e \frac{dt}{t} = 1.$$

( $e = 2.718281\dots$ ).



# Natural Logarithms (Sect. 7.2)

- ▶ Definition as an integral.
- ▶ **The derivative and properties.**
- ▶ The graph of the natural logarithm.
- ▶ Integrals involving logarithms.
- ▶ Logarithmic differentiation.

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t}$$

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \quad \Rightarrow \quad \ln'(x) = \frac{1}{x}.$$



# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \quad \Rightarrow \quad \ln'(x) = \frac{1}{x}.$$



## Theorem (Chain rule)

*For every differentiable function  $u$  holds  $[\ln(u)]' = \frac{u'}{u}$ .*

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \Rightarrow \ln'(x) = \frac{1}{x}.$$



## Theorem (Chain rule)

*For every differentiable function  $u$  holds  $[\ln(u)]' = \frac{u'}{u}$ .*

Proof:

$$\frac{d \ln(u)}{dx} =$$

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \Rightarrow \ln'(x) = \frac{1}{x}.$$



## Theorem (Chain rule)

*For every differentiable function  $u$  holds  $[\ln(u)]' = \frac{u'}{u}$ .*

Proof:

$$\frac{d \ln(u)}{dx} = \frac{d \ln}{du}(u) \frac{du}{dx}$$



# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \Rightarrow \ln'(x) = \frac{1}{x}.$$



## Theorem (Chain rule)

*For every differentiable function  $u$  holds  $[\ln(u)]' = \frac{u'}{u}$ .*

Proof:

$$\frac{d \ln(u)}{dx} = \frac{d \ln}{du}(u) \frac{du}{dx} = \frac{1}{u} u'$$

# The derivative and properties

## Theorem (Derivative of $\ln$ )

*The Fundamental Theorem of Calculus implies  $\ln'(x) = \frac{1}{x}$ .*

Proof:

$$\ln(x) = \int_1^x \frac{dt}{t} \Rightarrow \ln'(x) = \frac{1}{x}.$$



## Theorem (Chain rule)

*For every differentiable function  $u$  holds  $[\ln(u)]' = \frac{u'}{u}$ .*

Proof:

$$\frac{d \ln(u)}{dx} = \frac{d \ln}{du}(u) \frac{du}{dx} = \frac{1}{u} u' \Rightarrow \frac{d \ln(u)}{dx}(x) = \frac{u'(x)}{u(x)}.$$



# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3)$$

# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3) = \frac{1}{x}$$

# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3) = \frac{1}{x} \quad \Rightarrow \quad y'(x) = \frac{1}{x}.$$

# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3) = \frac{1}{x} \quad \Rightarrow \quad y'(x) = \frac{1}{x}.$$

We also use chain rule,

$$z'(x) = \frac{1}{(2x^2 + \cos(x))} (4x - \sin(x))$$

# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3) = \frac{1}{x} \quad \Rightarrow \quad y'(x) = \frac{1}{x}.$$

We also use chain rule,

$$z'(x) = \frac{1}{(2x^2 + \cos(x))} (4x - \sin(x))$$

$$z'(x) = \frac{4x - \sin(x)}{2x^2 + \cos(x)}.$$





# The derivative and properties

## Example

Find the derivative of  $y(x) = \ln(3x)$ , and  $z(x) = \ln(2x^2 + \cos(x))$ .

**Solution:** We use the chain rule.

$$y'(x) = \frac{1}{(3x)} (3) = \frac{1}{x} \quad \Rightarrow \quad y'(x) = \frac{1}{x}.$$

We also use chain rule,

$$z'(x) = \frac{1}{(2x^2 + \cos(x))} (4x - \sin(x))$$

$$z'(x) = \frac{4x - \sin(x)}{2x^2 + \cos(x)}.$$



**Remark:**  $y(x) = \ln(3x)$ , satisfies  $y'(x) = \ln'(x)$ .

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a$

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x}$

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

Therefore  $\ln(ax) = \ln(x) + c$ .

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

Therefore  $\ln(ax) = \ln(x) + c$ . Evaluating at  $x = 1$  we obtain  $c$ .



# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

Therefore  $\ln(ax) = \ln(x) + c$ . Evaluating at  $x = 1$  we obtain  $c$ .

$$\ln(a) = \ln(1) + c$$

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

Therefore  $\ln(ax) = \ln(x) + c$ . Evaluating at  $x = 1$  we obtain  $c$ .

$$\ln(a) = \ln(1) + c \Rightarrow c = \ln(a)$$

# The derivative and properties

## Theorem (Algebraic properties)

*For every positive real numbers  $a$  and  $b$  holds,*

(a)  $\ln(ab) = \ln(a) + \ln(b)$ , (product rule);

(b)  $\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$ , (quotient rule);

(c)  $\ln\left(\frac{1}{a}\right) = -\ln(a)$ , (reciprocal rule);

(d)  $\ln(a^b) = b\ln(a)$ , (power rule).

Proof of (a): (only)

The function  $y(x) = \ln(ax)$  satisfies  $y'(x) = \frac{1}{ax} a = \frac{1}{x} = \ln'(x)$

Therefore  $\ln(ax) = \ln(x) + c$ . Evaluating at  $x = 1$  we obtain  $c$ .

$$\ln(a) = \ln(1) + c \Rightarrow c = \ln(a) \Rightarrow \ln(ax) = \ln(x) + \ln(a).$$

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

$$y = \ln[(x+1)^2] - \ln[3(x+2)],$$

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

$$y = \ln[(x+1)^2] - \ln[3(x+2)],$$

$$y = 2 \ln(x+1) - [\ln(3) + \ln(x+2)].$$

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

$$y = \ln[(x+1)^2] - \ln[3(x+2)],$$

$$y = 2 \ln(x+1) - [\ln(3) + \ln(x+2)].$$

The derivative of function  $y$  is:  $y' = 2 \frac{1}{(x+1)} - \frac{1}{(x+2)}.$



# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

$$y = \ln[(x+1)^2] - \ln[3(x+2)],$$

$$y = 2 \ln(x+1) - [\ln(3) + \ln(x+2)].$$

The derivative of function  $y$  is:  $y' = 2 \frac{1}{(x+1)} - \frac{1}{(x+2)}.$

$$y' = \frac{2(x+2) - (x+1)}{(x+1)(x+2)}$$

# The derivative and properties

## Example

Compute the derivative of  $y(x) = \ln \left[ \frac{(x+1)^2}{3(x+2)} \right]$ .

**Solution:** Before computing the derivative of  $y$ , we simplify it,

$$y = \ln[(x+1)^2] - \ln[3(x+2)],$$

$$y = 2 \ln(x+1) - [\ln(3) + \ln(x+2)].$$

The derivative of function  $y$  is:  $y' = 2 \frac{1}{(x+1)} - \frac{1}{(x+2)}.$

$$y' = \frac{2(x+2) - (x+1)}{(x+1)(x+2)} \Rightarrow y' = \frac{(x+3)}{(x+1)(x+2)}. \quad \triangleleft$$

# Natural Logarithms (Sect. 7.2)


- ▶ Definition as an integral.
- ▶ The derivative and properties.
- ▶ The graph of the natural logarithm.
- ▶ Integrals involving logarithms.
- ▶ Logarithmic differentiation.

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.




# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.




Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .


(a): If  $x = e^n$ ,

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .


(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e)$

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ .




# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence


$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence

$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$


(b): If  $x = \frac{1}{e^n}$ ,

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence

$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$


(b): If  $x = \frac{1}{e^n}$ , then  $\ln\left(\frac{1}{e^n}\right) = -\ln(e^n)$

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence

$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$


(b): If  $x = \frac{1}{e^n}$ , then  $\ln\left(\frac{1}{e^n}\right) = -\ln(e^n) = -n \ln(e)$

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence

$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$


(b): If  $x = \frac{1}{e^n}$ , then  $\ln\left(\frac{1}{e^n}\right) = -\ln(e^n) = -n \ln(e) = -n$ .

# The graph of the natural logarithm

## Remarks:

The graph of  $\ln$  function has:

- (a) A vertical asymptote at  $x = 0$ .
- (b) No horizontal asymptote.



Proof: Recall  $e = 2.718281... > 1$  and  $\ln(e) = 1$ .

(a): If  $x = e^n$ , then  $\ln(e^n) = n \ln(e) = n$ . Hence

$$\lim_{x \rightarrow \infty} \ln(x) = \infty.$$

(b): If  $x = \frac{1}{e^n}$ , then  $\ln\left(\frac{1}{e^n}\right) = -\ln(e^n) = -n \ln(e) = -n$ . Hence

$$\lim_{x \rightarrow 0^+} \ln(x) = -\infty.$$

# Natural Logarithms (Sect. 7.2)

- ▶ Definition as an integral.
- ▶ The derivative and properties.
- ▶ The graph of the natural logarithm.
- ▶ **Integrals involving logarithms.**
- ▶ Logarithmic differentiation.

## Integrals involving logarithms.

Remark: It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .



## Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

# Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

And for  $x < 0$ , we have that  $-x > 0$ ,

## Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

And for  $x < 0$ , we have that  $-x > 0$ , then,

$$\int \frac{dx}{x} = \int \frac{(-dx)}{(-x)}$$

## Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

And for  $x < 0$ , we have that  $-x > 0$ , then,

$$\int \frac{dx}{x} = \int \frac{(-dx)}{(-x)} = \ln(-x) + c, \quad -x > 0.$$

# Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

And for  $x < 0$ , we have that  $-x > 0$ , then,

$$\int \frac{dx}{x} = \int \frac{(-dx)}{(-x)} = \ln(-x) + c, \quad -x > 0.$$

We conclude,

$$\int \frac{dx}{x} = \begin{cases} \ln(-x) + c & \text{if } x < 0, \\ \ln(x) + c & \text{if } x > 0. \end{cases}$$

# Integrals involving logarithms.

**Remark:** It holds  $\int \frac{dx}{x} = \ln(|x|) + c$  for  $x \neq 0$  and  $c \in \mathbb{R}$ .

Indeed, for  $x > 0$  this is the definition of logarithm.

And for  $x < 0$ , we have that  $-x > 0$ , then,

$$\int \frac{dx}{x} = \int \frac{(-dx)}{(-x)} = \ln(-x) + c, \quad -x > 0.$$

We conclude,

$$\int \frac{dx}{x} = \begin{cases} \ln(-x) + c & \text{if } x < 0, \\ \ln(x) + c & \text{if } x > 0. \end{cases}$$

**Remark:** It also holds  $\int \frac{f'(x)}{f(x)} dx = \ln(|f(x)|) + c$ , for  $f(x) \neq 0$ .

# Integrals involving logarithms.

Remarks:

$$(a) \int \tan(x) dx = -\ln(|\cos(x)|) + c.$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) \, dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) \, dx = \int \frac{\sin(x)}{\cos(x)} \, dx$$



# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) \, dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) \, dx = \int \frac{\sin(x)}{\cos(x)} \, dx \quad u = \cos(x),$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u}$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ .

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx \quad u = \sin(x),$$



# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx \quad u = \sin(x), \quad du = \cos(x) dx.$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx \quad u = \sin(x), \quad du = \cos(x) dx.$$

$$\int \cot(x) dx = \int \frac{du}{u}$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx \quad u = \sin(x), \quad du = \cos(x) dx.$$

$$\int \cot(x) dx = \int \frac{du}{u} = \ln(|u|) + c$$

# Integrals involving logarithms.

Remarks:

(a)  $\int \tan(x) dx = -\ln(|\cos(x)|) + c$ . Indeed,

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx \quad u = \cos(x), \quad du = -\sin(x) dx.$$

$$\int \tan(x) dx = - \int \frac{du}{u} = -\ln(|u|) + c = -\ln(|\cos(x)|) + c.$$

(b)  $\int \cot(x) dx = \ln(|\sin(x)|) + c$ . Indeed,

$$\int \cot(x) dx = \int \frac{\cos(x)}{\sin(x)} dx \quad u = \sin(x), \quad du = \cos(x) dx.$$

$$\int \cot(x) dx = \int \frac{du}{u} = \ln(|u|) + c = \ln(|\sin(x)|) + c.$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt,$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t),$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t), \quad du = -\sin(t) dt.$$



# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t), \quad du = -\sin(t) dt.$$

$$y(t) = \int \frac{3(-du)}{u}$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t), \quad du = -\sin(t) dt.$$

$$y(t) = \int \frac{3(-du)}{u} = -3 \int \frac{du}{u}$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t), \quad du = -\sin(t) dt.$$

$$y(t) = \int \frac{3(-du)}{u} = -3 \int \frac{du}{u} = -3 \ln(|u|) + c$$

# Integrals involving logarithms.

## Example

Find  $y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt.$

## Solution:

$$y(t) = \int \frac{3 \sin(t)}{(2 + \cos(t))} dt, \quad u = 2 + \cos(t), \quad du = -\sin(t) dt.$$

$$y(t) = \int \frac{3(-du)}{u} = -3 \int \frac{du}{u} = -3 \ln(|u|) + c$$

We conclude that  $y(t) = -3 \ln(|2 + \cos(t)|) + c.$



# Natural Logarithms (Sect. 7.2)

- ▶ Definition as an integral.
- ▶ The derivative and properties.
- ▶ The graph of the natural logarithm.
- ▶ Integrals involving logarithms.
- ▶ **Logarithmic differentiation.**

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

**Example**

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** First compute  $\ln[y(x)]$



# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** First compute  $\ln[y(x)] = \ln\left[\frac{x^3(x+2)^2}{\cos^3(x)}\right]$ ,

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** First compute  $\ln[y(x)] = \ln\left[\frac{x^3(x+2)^2}{\cos^3(x)}\right]$ ,

$$\ln[y(x)] = \ln[x^3(x+2)^2] - \ln[\cos^3(x)],$$

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** First compute  $\ln[y(x)] = \ln\left[\frac{x^3(x+2)^2}{\cos^3(x)}\right]$ ,

$$\ln[y(x)] = \ln[x^3(x+2)^2] - \ln[\cos^3(x)],$$

$$\ln[y(x)] = \ln[x^3] + \ln[(x+2)^2] - \ln[\cos^3(x)],$$

# Logarithmic differentiation

**Remark:** Logarithms can be used to simplify the derivative of complicated functions.

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** First compute  $\ln[y(x)] = \ln\left[\frac{x^3(x+2)^2}{\cos^3(x)}\right]$ ,

$$\ln[y(x)] = \ln[x^3(x+2)^2] - \ln[\cos^3(x)],$$

$$\ln[y(x)] = \ln[x^3] + \ln[(x+2)^2] - \ln[\cos^3(x)],$$

$$\ln[y(x)] = 3\ln(x) + 2\ln(x+2) - 3\ln[\cos(x)].$$

# Logarithmic differentiation

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

Solution: Recall:  $\ln[y(x)] = 3 \ln(x) + 2 \ln(x+2) - 3 \ln[\cos(x)]$ .

# Logarithmic differentiation

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

Solution: Recall:  $\ln[y(x)] = 3\ln(x) + 2\ln(x+2) - 3\ln[\cos(x)]$ .

$$\frac{y'(x)}{y(x)} = \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)}.$$

# Logarithmic differentiation

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

Solution: Recall:  $\ln[y(x)] = 3\ln(x) + 2\ln(x+2) - 3\ln[\cos(x)]$ .

$$\frac{y'(x)}{y(x)} = \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)}.$$

$$y'(x) = \left[ \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)} \right] y(x).$$

# Logarithmic differentiation

## Example

Find the derivative of  $y(x) = \frac{x^3(x+2)^2}{\cos^3(x)}$ .

**Solution:** Recall:  $\ln[y(x)] = 3\ln(x) + 2\ln(x+2) - 3\ln[\cos(x)]$ .

$$\frac{y'(x)}{y(x)} = \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)}.$$

$$y'(x) = \left[ \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)} \right] y(x).$$

We conclude that

$$y'(x) = \left[ \frac{3}{x} + \frac{2}{(x+2)} + \frac{3\sin(x)}{\cos(x)} \right] \frac{\cos^3(x)}{x^3(x+2)^2}. \quad \triangleleft$$



# The inverse function (Sect. 7.1)

- ▶ One-to-one functions.
- ▶ The inverse function
- ▶ The graph of the inverse function.
- ▶ Derivatives of the inverse function.

# One-to-one functions

Remark:

- ▶ Not every function is invertible.

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \quad \Rightarrow \quad f(x_1) \neq f(x_2).$$

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \quad \Rightarrow \quad f(x_1) \neq f(x_2).$$

## Example

Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

### Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

### Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .
4.  $y = \sin(x)$ ,  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .



# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .
4.  $y = \sin(x)$ ,  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .

Not Invertible:

1.  $y = x^2$ , for  $x \in [-a, a]$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

### Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .
4.  $y = \sin(x)$ ,  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .

### Not Invertible:

1.  $y = x^2$ , for  $x \in [-a, a]$ .
2.  $y = |x|$ , for  $x \in [-a, a]$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

### Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .
4.  $y = \sin(x)$ ,  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .

### Not Invertible:

1.  $y = x^2$ , for  $x \in [-a, a]$ .
2.  $y = |x|$ , for  $x \in [-a, a]$ .
3.  $y = \cos(x)$ ,  $x \in [-a, a]$ .

# One-to-one functions

## Remark:

- ▶ Not every function is invertible.
- ▶ Only one-to-one functions are invertible.

## Definition

A function  $f : D \rightarrow \mathbb{R}$  is called *one-to-one* (injective) iff for every  $x_1, x_2 \in D$  holds

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2).$$

## Example

### Invertible:

1.  $y = x^3$ , for  $x \in \mathbb{R}$ .
2.  $y = x^2$ , for  $x \in [0, b]$ .
3.  $y = \sqrt{x}$ , for  $x \in [0, \infty)$ .
4.  $y = \sin(x)$ ,  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .

### Not Invertible:

1.  $y = x^2$ , for  $x \in [-a, a]$ .
2.  $y = |x|$ , for  $x \in [-a, a]$ .
3.  $y = \cos(x)$ ,  $x \in [-a, a]$ .
4.  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1$$



# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ ,

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

## Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta)$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta)$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ ,



# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ , and taking  $x_1 = \pi/4$ ,  $x_2 = 3\pi/4$ ,

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ , and taking  $x_1 = \pi/4$ ,  $x_2 = 3\pi/4$ ,

$$f(\pi/4) = \sin(\pi/4)$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ , and taking  $x_1 = \pi/4$ ,  $x_2 = 3\pi/4$ ,

$$f(\pi/4) = \sin(\pi/4) = \sin(\pi - \pi/4)$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ , and taking  $x_1 = \pi/4$ ,  $x_2 = 3\pi/4$ ,

$$f(\pi/4) = \sin(\pi/4) = \sin(\pi - \pi/4) = \sin(3\pi/4)$$

# One-to-one functions

## Example

Verify that the functions below are not one-to-one:

(a)  $y = x^2$ , for  $x \in [-a, a]$ .

(b)  $y = \cos(x)$ ,  $x \in [-a, a]$ .

(c)  $y = \sin(x)$ , for  $x \in [0, \pi]$ .

Solution:

(a) For  $x_1 = -1$ ,  $x_2 = 1$  we have that  $x_1 \neq x_2$  and

$$f(x_1) = (-1)^2 = 1 = 1^2 = f(x_2).$$

(b) Recalling that  $\cos(\theta) = \cos(-\theta)$ , and taking  $x_1 = -\theta$ ,  $x_2 = \theta$ ,

$$f(-\theta) = \cos(-\theta) = \cos(\theta) = f(\theta).$$

(c) Since  $\sin(\theta) = \sin(\pi - \theta)$ , and taking  $x_1 = \pi/4$ ,  $x_2 = 3\pi/4$ ,

$$f(\pi/4) = \sin(\pi/4) = \sin(\pi - \pi/4) = \sin(3\pi/4) = f(3\pi/4). \triangleleft$$

# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ ,



# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ ,

# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ , that means  $y_0 = f(x_1) = f(x_2)$ .

# One-to-one functions

**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ , that means  $y_0 = f(x_1) = f(x_2)$ . Hence  $f$  is not one-to-one. □

# One-to-one functions


**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ , that means  $y_0 = f(x_1) = f(x_2)$ . Hence  $f$  is not one-to-one. □

## Example



# One-to-one functions



**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ , that means  $y_0 = f(x_1) = f(x_2)$ . Hence  $f$  is not one-to-one. □

## Example



# One-to-one functions




**Remark:** By looking at the graph of the function one can determine whether the function is one-to-one or not.

## Theorem (Horizontal line test)

*A function  $f : D \rightarrow R$  is one-to-one iff the function graph intersects every horizontal line at most once.*

**Proof:** If a function  $f$  intersects the horizontal line  $y = y_0$  at  $(x_1, f(x_1))$  and  $(x_2, f(x_2))$ , with  $x_1 \neq x_2$ , that means  $y_0 = f(x_1) = f(x_2)$ . Hence  $f$  is not one-to-one. □

## Example



# The inverse function (Sect. 7.1)

- ▶ One-to-one functions.
- ▶ **The inverse function**
- ▶ The graph of the inverse function.
- ▶ Derivatives of the inverse function.

# The inverse function

**Remark:** Only one-to-one functions are invertible.



# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ ,

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ , that is,  $y = 2x - 3$ .

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ , that is,  $y = 2x - 3$ . Find  $x$  in the expression above,

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ , that is,  $y = 2x - 3$ . Find  $x$  in the expression above,

$$2x = y + 3$$

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ , that is,  $y = 2x - 3$ . Find  $x$  in the expression above,

$$2x = y + 3 \quad \Rightarrow \quad x = \frac{1}{2}y + \frac{3}{2}.$$

# The inverse function

**Remark:** Only one-to-one functions are invertible.

## Definition

The *inverse* of a one-to-one function  $f : D \rightarrow R$  is the function  $f^{-1} : R \rightarrow D$  defined for all  $x \in D$  and all  $y \in R$  as follows

$$f^{-1}(y) = x \quad \Leftrightarrow \quad y = f(x).$$

## Example

Find the inverse of  $f(x) = 2x - 3$ .

**Solution:** Denote  $y = f(x)$ , that is,  $y = 2x - 3$ . Find  $x$  in the expression above,

$$2x = y + 3 \quad \Rightarrow \quad x = \frac{1}{2}y + \frac{3}{2}.$$

Then, the inverse function is  $f^{-1}(y) = \frac{1}{2}y + \frac{3}{2}$ .





# The inverse function

Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

# The inverse function

Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ .

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x))$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3)$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3]$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3] = x.$$



# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3] = x.$$

$$f(f^{-1}(y))$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3] = x.$$

$$f(f^{-1}(y)) = f\left(\frac{1}{2}(y + 3)\right)$$

# The inverse function

## Remark:

- If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3] = x.$$

$$f(f^{-1}(y)) = f\left(\frac{1}{2}(y + 3)\right) = 2\left[\frac{1}{2}(y + 3)\right] - 3$$

# The inverse function

## Remark:

- ▶ If  $f^{-1}$  is the inverse of  $f$ , then holds

$$(f^{-1} \circ f)(x) = x, \quad (f \circ f^{-1})(y) = y.$$

- ▶ Equivalently,

$$f^{-1}(f(x)) = x, \quad f(f^{-1}(y)) = y.$$

## Example

Verify the relations above for  $f(x) = 2x - 3$ .

**Solution:** Recall:  $f^{-1}(y) = (y + 3)/2$ . Hence

$$f^{-1}(f(x)) = f^{-1}(2x - 3) = \frac{1}{2}[(2x - 3) + 3] = x.$$

$$f(f^{-1}(y)) = f\left(\frac{1}{2}(y + 3)\right) = 2\left[\frac{1}{2}(y + 3)\right] - 3 = y. \quad \triangleleft$$

# The inverse function (Sect. 7.1)

- ▶ One-to-one functions.
- ▶ The inverse function
- ▶ **The graph of the inverse function.**
- ▶ Derivatives of the inverse function.


# The graph of the inverse function

**Remark:** The graph of the function  $f^{-1}$  is obtained reflecting the graph of  $f$  along the line  $y = x$ .

# The graph of the inverse function

**Remark:** The graph of the function  $f^{-1}$  is obtained reflecting the graph of  $f$  along the line  $y = x$ .



## Example



# The graph of the inverse function

**Remark:** The graph of the function  $f^{-1}$  is obtained reflecting the graph of  $f$  along the line  $y = x$ .

## Example






# The graph of the inverse function

**Remark:** The graph of the function  $f^{-1}$  is obtained reflecting the graph of  $f$  along the line  $y = x$ .

## Example



# The inverse function (Sect. 7.1)


- ▶ One-to-one functions.
- ▶ The inverse function
- ▶ The graph of the inverse function.
- ▶ **Derivatives of the inverse function.**

# Derivatives of the inverse function.

**Remark:** The derivative values of a function and its inverse are deeply related.



# Derivatives of the inverse function.

**Remark:** The derivative values of a function and its inverse are deeply related.





# Derivatives of the inverse function.

**Remark:** The derivative values of a function and its inverse are deeply related.



# Derivatives of the inverse function.

**Remark:** The derivative values of a function and its inverse are deeply related.





## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable.*

# Derivatives of the inverse function.

**Remark:** The derivative values of a function and its inverse are deeply related.



## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$



# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

## Example

Verify the Theorem above for  $f(x) = 2x - 3$ .

# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

## Example

Verify the Theorem above for  $f(x) = 2x - 3$ .

**Solution:** This case is simple because  $f'(x) = 2$ , constant.

# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

## Example

Verify the Theorem above for  $f(x) = 2x - 3$ .

**Solution:** This case is simple because  $f'(x) = 2$ , constant.

Since  $f^{-1}(y) = \frac{1}{2}(y + 3)$ ,

# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

## Example

Verify the Theorem above for  $f(x) = 2x - 3$ .

**Solution:** This case is simple because  $f'(x) = 2$ , constant.

Since  $f^{-1}(y) = \frac{1}{2}(y + 3)$ , then  $(f^{-1})'(y) = \frac{1}{2}$ , constant.

# Derivatives of the inverse function.

## Theorem (Derivative for inverse functions)

*If the invertible function  $f : D \rightarrow R$  is differentiable and  $f'(x) \neq 0$  for every  $x \in D$ , then the function  $f^{-1} : R \rightarrow D$  is differentiable. Furthermore, for every  $y \in R$  holds*

$$(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$$

## Example

Verify the Theorem above for  $f(x) = 2x - 3$ .

**Solution:** This case is simple because  $f'(x) = 2$ , constant.

Since  $f^{-1}(y) = \frac{1}{2}(y + 3)$ , then  $(f^{-1})'(y) = \frac{1}{2}$ , constant.

Therefore,  $(f^{-1})' = \frac{1}{f'}$ .



# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function.

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ ,



# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3}$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3}$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

$$f'(x) = 3x^2$$



# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

$$f'(x) = 3x^2 \quad \Rightarrow \quad f'(f^{-1}(y)) = 3(f^{-1}(y))^2$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

$$f'(x) = 3x^2 \quad \Rightarrow \quad f'(f^{-1}(y)) = 3(f^{-1}(y))^2 = 3(y^{1/3})^2.$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \quad \Rightarrow \quad x = y^{1/3} \quad \Rightarrow \quad f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

$$f'(x) = 3x^2 \quad \Rightarrow \quad f'(f^{-1}(y)) = 3(f^{-1}(y))^2 = 3(y^{1/3})^2.$$

We conclude that  $\frac{1}{3y^{2/3}} = (f^{-1})'(y)$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3$ .

## Solution:

We first compute the inverse function. Denote  $y = f(x)$ , then

$$y = x^3 \Rightarrow x = y^{1/3} \Rightarrow f^{-1}(y) = y^{1/3}.$$

Compute now the derivative of the inverse function,

$$(f^{-1})'(y) = \frac{1}{3} y^{-2/3} = \frac{1}{3} \frac{1}{y^{2/3}}.$$

Compute now the derivative of the original function,

$$f'(x) = 3x^2 \Rightarrow f'(f^{-1}(y)) = 3(f^{-1}(y))^2 = 3(y^{1/3})^2.$$

We conclude that  $\frac{1}{3y^{2/3}} = (f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}.$



# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

## Derivatives of the inverse function.

### Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ ,

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .



# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

## Derivatives of the inverse function.

### Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

Then the formula in the Theorem,  $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$ ,

## Derivatives of the inverse function.

### Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

Then the formula in the Theorem,  $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$ ,

for  $y = 9$  implies

$$(f^{-1})'(9) = \frac{1}{f'(f^{-1}(9))}$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

Then the formula in the Theorem,  $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$ ,

for  $y = 9$  implies

$$(f^{-1})'(9) = \frac{1}{f'(f^{-1}(9))} = \frac{1}{f'(2)}$$

# Derivatives of the inverse function.

## Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

Then the formula in the Theorem,  $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$ ,

for  $y = 9$  implies

$$(f^{-1})'(9) = \frac{1}{f'(f^{-1}(9))} = \frac{1}{f'(2)} = \frac{1}{3(4) + 1}.$$

## Derivatives of the inverse function.

### Example

Verify the Theorem above for  $f(x) = x^3 + x - 1$  at  $x = 2$ .

**Solution:** In this case is difficult to find  $f^{-1}(y)$  for  $y \in \mathbb{R}$ .

But at  $x = 2$  we have  $f(2) = 2^3 + 2 - 1$ , that is,  $f(2) = 9$ .

Therefore  $f^{-1}(9) = 2$ .

Compute now the derivative of the original function,

$$f'(x) = 3x^2 + 1.$$

Then the formula in the Theorem,  $(f^{-1})'(y) = \frac{1}{f'(f^{-1}(y))}$ ,

for  $y = 9$  implies

$$(f^{-1})'(9) = \frac{1}{f'(f^{-1}(9))} = \frac{1}{f'(2)} = \frac{1}{3(4) + 1}.$$

We conclude that  $(f^{-1})'(y) = \frac{1}{13}$ .

